



Clinical Trial Details (PDF Generation Date :- Sun, 01 Oct 2023 02:52:11 GMT)

CTRI Number	CTRI/2020/12/029735 [Registered on: 11/12/2020] - Trial Registered Prospectively		
Last Modified On	13/05/2021		
Post Graduate Thesis	No		
Type of Trial	Interventional		
Type of Study	Ayurveda		
Study Design	Randomized, Parallel Group Trial		
Public Title of Study	A Study to Assess the Safety and Efficacy of CoroQuil-Zn 750 in Comparison to the Standard of Care for the Treatment of Mild to Moderate COVID-19.		
Scientific Title of Study	An Open-Label, Two-Arm, Parallel Design, Prospective, Single-Center, Phase 3 study to assess the safety and efficacy of CoroQuil-Zn in comparison to the standard of care for the treatment of mild to moderate COVID-19		
Secondary IDs if Any	Secondary ID	Identifier	
	NIL	NIL	
Details of Principal Investigator or overall Trial Coordinator (multi-center study)	Details of Principal Investigator		
	Name	Dr Sambashiva AC	
	Designation	Principal Investigator	
	Affiliation	Sri Lakshmi Super speciality hospital	
	Address	Department of General Medicine, No. 5,6,7, 1st Cross, Kaggadasapura Main Rd, Nagappareddy Layout, C V Raman Nagar, Bengaluru, Karnataka Bangalore KARNATAKA 560093 India	
	Phone	6364147989	
	Fax		
	Email	info@samahitha.com	
	Details Contact Person (Scientific Query)	Details Contact Person (Scientific Query)	
		Name	Mr Srivatsa
Designation		Project Manager & MD	
Affiliation		Samahitha Research Solutions	
Address		#1278, 25th main road, 40th Cross Road, 9th Block, Jayanagar, Bengaluru Bangalore KARNATAKA 560069 India	
Phone		9742797117	
Fax			
Email		sri@samahitha.com	
Details Contact Person (Public Query)	Details Contact Person (Public Query)		
	Name	Disha Shetty	
	Designation	Incharge Clinical Operations	
	Affiliation	Samahitha Research Solutions	
	Address	#1278, 25th main road, 40th Cross Road, 9th Block, Jayanagar, Bengaluru Bangalore KARNATAKA 560069	



	India			
Phone	6364147979			
Fax				
Email	disha@samahitha.com			
Source of Monetary or Material Support	Source of Monetary or Material Support			
	> Remedium Therapeutics Private Limited			
Primary Sponsor	Primary Sponsor Details			
Name	Remedium Therapeutics Private Limited			
Address	No.1320 2nd floor 33rd street 7th sector KK Nagar Chennai Tamil Nadu			
Type of Sponsor	Pharmaceutical industry-Indian			
Details of Secondary Sponsor	Name	Address		
	Remedium Therapeutics Private Limited	No.1320 2nd floor 33rd street 7th sector KK Nagar Chennai Tamil Nadu		
Countries of Recruitment	List of Countries			
	India			
Sites of Study	Name of Principal Investigator	Name of Site	Site Address	Phone/Fax/Email
	Dr Sambashiva AC	Sri Lakshmi Super Speciality Hospital	Sri Lakshmi super speciality hospital, Department of General Medicine No. 5,6,7, 1st cross, Kaggadasapura main Rd, nagappareddy layout, C V Raman Nagar, Bengaluru, Karnataka 560093 Bangalore KARNATAKA	6364147979 info@samahitha.com
Details of Ethics Committee	Name of Committee	Approval Status	Date of Approval	Is Independent Ethics Committee?
	PRANAV DIABETES CENTER ETHICS COMMITTEE	Approved	03/12/2020	No
Regulatory Clearance Status from DCGI	Status	Date		
	Not Applicable	No Date Specified		
Health Condition / Problems Studied	Health Type	Condition		
	Patients	Coronavirus as the cause of diseases classified elsewhere		
Intervention / Comparator Agent	Type	Name	Details	
	Intervention	CoroQuil-Zn 750mg	750mg x 2, administered orally three times a day for 14 days	
	Comparator Agent	COVID Standard Care	As per the sites COVID SOP	
Inclusion Criteria	Inclusion Criteria			
	Age From	18.00 Year(s)		
	Age To	75.00 Year(s)		
	Gender	Both		
	Details	1. Written signed and dated informed consent (patient or LAR). 2. Both genders, aged ?18 to ?75 years, 3. Patients with RT-qPCR confirmed COVID-19 patients 4. SpO2?90% and		



	respiratory rate \geq 30/minute 5. Healthy adult patients with ASA I to II 6. A score of between 3 to 5 on the Modified WHO Ordinal Scale for Clinical Improvement (refer protocol appendix 23.1) 7. Patients who agree to abide by the study requirements				
Exclusion Criteria	Exclusion Criteria				
Details	<ol style="list-style-type: none"> 1. Pregnant and lactating women 2. Children \leq 75 years 3. SpO₂ \leq 93% / minute 4. Patients having persistent nausea/vomiting 5. Need for direct admission to the intensive care unit for mechanical ventilation 6. Underlying chronic obstructive pulmonary disease stage III-IV 7. Patients simultaneously participating in another clinical study. 8. History of stroke with significant neurologic deficit. 9. Patients with any concurrent pre-existing severe/uncontrolled, clinically significant systemic disease [e.g. heart failure (NYHA 2 or above)], cancer, liver disease, kidney disease or anaemia etc.) that, in the opinion of investigator precludes the subject's participation in the study or interferes with the interpretation of the study results. 10. Patients with history of serology tests positive for hepatitis B, hepatitis C, or human immunodeficiency virus. 11. Medical or psychological conditions deemed by the investigators to interfere with successful participation in the study 12. A subject who is judged by the investigator as inappropriate to participate in the study for any reason other than those mentioned above. 				
Method of Generating Random Sequence	Coin toss, Lottery, toss of dice, shuffling cards etc				
Method of Concealment	Not Applicable				
Blinding/Masking	Open Label				
Primary Outcome	<table border="1"> <thead> <tr> <th>Outcome</th> <th>Timepoints</th> </tr> </thead> <tbody> <tr> <td>Negative result for qRT-PCR testing for COVID</td> <td>14 days</td> </tr> </tbody> </table>	Outcome	Timepoints	Negative result for qRT-PCR testing for COVID	14 days
Outcome	Timepoints				
Negative result for qRT-PCR testing for COVID	14 days				
Secondary Outcome	<table border="1"> <thead> <tr> <th>Outcome</th> <th>Timepoints</th> </tr> </thead> <tbody> <tr> <td>percentage of patients showing negative qRT-PCR earlier than the End of study visit</td> <td>day 5, day 7</td> </tr> </tbody> </table>	Outcome	Timepoints	percentage of patients showing negative qRT-PCR earlier than the End of study visit	day 5, day 7
Outcome	Timepoints				
percentage of patients showing negative qRT-PCR earlier than the End of study visit	day 5, day 7				
Target Sample Size	Total Sample Size=100 Sample Size from India=100 Final Enrollment numbers achieved (Total)=114 Final Enrollment numbers achieved (India)=114				
Phase of Trial	Phase 3				
Date of First Enrollment (India)	15/12/2020				
Date of First Enrollment (Global)	No Date Specified				
Estimated Duration of Trial	Years=0 Months=4 Days=0				
Recruitment Status of Trial (Global)	Not Applicable				
Recruitment Status of Trial (India)	Completed				
Publication Details	NIL				
Brief Summary	This was an open-label, two-arm, parallel design, prospective, single-centre, phase 3 study to				



assess the safety and efficacy of CoroQuil-Zn when compared to the Standard of Care [Mild cases: Azithromycin 500mg tab (once daily) + Corticosteroids (Twice daily) + Paracetamol (Twice daily) + Aspirin 500mg (Twice daily); Moderate Cases: Piperceillin tazobactam 4.5mg inj. or Ceftriaxone 1gm inj.(Once daily) + Corticosteroid (Twice daily) + Enoxiparin 40 (Twice daily) + Paracetamol 650mg (Twice daily); IV infusion if required (only in case of weakness); All Antibiotics for 6 days and other medications based on the reports and PI discretion] for the treatment of mild to moderate COVID-19 infection.

- [120 patients screened and 114 randomized in this study.](#)
- Both inpatients and outpatients were enrolled in this study.
- 57% males and 42% females were enrolled in this study.
- The median age group was 31 years in this study.
- The study drug was well-tolerated by the subjects and no adverse drug reactions were observed.
- There were no deaths reported in both the treatment groups.
- Clinical Response & Clinical Cure at the Day 7 visit: 98% of the patients in the CoroQuil-Zn group showed a clinical response when compared to the Standard of Care group (82%) which is statistically significant (p-value: 0.0035).
- The median time to Clinical Response & Clinical Cure was 3 days in the CoroQuil-Zn group and 6 days in the Standard of Care group which is statistically significant (p-value: <0.0001).
- Virological Cure at Day 7 visit: Majority of the patients in the CoroQuil-Zn arm (95%) and showed virological cure as compared to the Standard of Care group (59%) which is statistically significant (p-value: <0.0001).
- Virological Cure at Day 14 visit: Majority (97%) of the patients in the CoroQuil-Zn group showed virological cure when compared to the Standard of Care group (86%) which is statistically significant (p-value: 0.0372).
- The median time for Virological Cure was 6 days for the CoroQuil-Zn group and 7 days for the Standard of Care group which is a statistically significant (p-value: <0.0001).
- The severity of Dyspnea at the Day 7 visit: There is a significant reduction in Dyspnoea with CoroQuil-Zn group as compared to the Standard of Care group which is statistically significant (p-value: <0.0001).
- The severity of Dyspnea at the Day 14 visit: There is no significant reduction in Dyspnea with the CoroQuil-Zn group as compared to the Standard of Care group on Day 14.



- Baseline IL-6 levels compared to Day 7 & Day 14 visit: There is a statistically significant reduction in IL-6 levels at day 7 (p-value: <0.0001) & Day 14 (p-value: <0.0001) in the CoroQuil-Zn group which is not seen in the Standard of Care group when compared to the baseline visit. There is statistically significant change seen between the two treatment groups at day 7 (p-value: 0.002) and day 14 visit (p-value: 0.0046) when compared to the baseline visit
- Baseline CRP, D-Dimer & Serum Ferritin levels compared to Day 7 & Day 14 visit: There is no statistically significant reduction observed between the two treatment groups at day 7 and day 14 visits when compared to the baseline visit.
- No clinically significant findings were observed in the physical examination, vitals, clinical laboratory tests, 12-lead ECG, chest X-Ray and routine urine analysis.
- A total of 05 adverse events (AEs) were reported in this study. All AEs were mild in severity, which resolved without sequelae.

Overall, at Day 7 visit in the CoroQuil-Zn treatment group, a statistically significant Clinical response, Clinical cure, Virological cure and reduction in the severity of Dyspnoea was observed.

Conclusion:

[During the primary analysis, a numerically higher positive trend was observed in the clinical response. Throughout the trial, a statistically significant improvement was seen in the CoroQuil-Zn treatment group for Clinical response, Clinical cure, and Virological cure, reduction in the severity of Dyspnoea and recovery rate in the secondary analysis. The median time to Clinical response, Clinical cure, and Virological cure was also statistically significant in the CoroQuil-Zn treatment group.](#)

From the analysis of this trial data, it is evident that capsule CoroQuil-Zn750 mg benefits mild to moderate COVID-19 patients. Administering CoroQuil-Zn 750mg will reduce the patient load on the hospital, recovery at home and may fulfil the current unmet need during this ongoing pandemic.

